arXiv:hep-ph/0206283v1 27 Jun 2002

CHARGED HIGGS IN MODELS WITH SINGLET NEUTRINO IN LARGE EXTRA DIMENSIONS [a]

A. DEANDREA
Institut de Physique Nucléaire, Université de Lyon I
4 rue E. Fermi, F-69622 Villeurbanne Cedex, France

The charged Higgs decay in the channel $H^- \to \tau_L^- \nu$ in models with a singlet neutrino can provide a test of large extra dimensions models with TeV scale quantum gravity since in the standard two Higgs doublet model type II, $H^- \to \tau_L^- \nu$ is suppressed. In the present study, we focus on the decay $H^- \to \tau_L^- \psi$ at the LHC for Higgs masses larger than the top-quark mass.

1 Introduction

Models with extra dimensions postulate the existence of δ additional spatial dimensions of size R where gravity and perhaps other fields freely propagate while the SM degrees of freedom are confined to (3+1)-dimensional wall (4D) of the higher dimensional space. The true scale of gravity, or fundamental Planck scale M_*, of the $(4+\delta)$D space time is related to the reduced 4D Planck scale M_{Pl}, by $M_{Pl}^2 = R^\delta M_*^{\delta+2}$, where $M_{Pl} = 2.4 \times 10^{18}$ GeV is related to the usual Planck mass 1.2×10^{19} GeV $= \sqrt{8\pi} M_{Pl}$. Since no experimental deviations from Newtonian gravity are observed at distances above 0.2 mm [1], the extra dimensions must be smaller or equal to the sub-millimeter scale with M_* as low as few TeV and $\delta \geq 2$.

The spectrum of many extensions of the SM includes a charged Higgs state. H^- decays to the right handed τ^- through the τ Yukawa coupling: $H^- \to \tau_R^- \bar{\nu}$. The H^- decay to left handed τ^- is completely suppressed in the Minimal Supersymmetric Standard Model (MSSM). However, in the scenario of singlet neutrino in large extra dimensions, H^- can decay to both right handed and left handed τ^- depending on the parameters M_*, m_D, δ, $m_{H^\pm}$ and $\tan\beta$: $H^- \to \tau_R^- \bar{\nu} + \tau_L^- \psi$, where ψ is a bulk neutrino and ν is dominantly a light neutrino with a small admixture of the Kaluza-Klein modes [2] of the order $mR/|n|$. The measurement of the polarisation asymmetry, can be used to distinguish between the ordinary two Higgs Doublet Model of type II (2HDM-II) and the scenario of singlet neutrino in large extra dimensions.

---

[a] based on work done in collaboration with K. Assamagan. Talk given at XXXVIIth Rencontres de Moriond, ElectroWeak Interactions and Unified Theories, March 9th - 16th 2002, Les Arcs, France

2 $H^\pm$ Production and Decays

The charged Higgs decay to right handed τ, $H^- \to \tau_R^- \bar{\nu}$ have been extensively studied for the LHC[3,4]. Here we discuss the possibility to observe $H^- \to \tau_L^- \psi$ at the LHC above the top-quark mass[5]. Table 1 shows the parameters selected for the current analysis. We assume a heavy SUSY spectrum with maximal mixing. We consider the $2 \to 2$ production process where the charged Higgs is produced with a top-quark, $gb \to tH^\pm$. Further, we require the hadronic decay of the top-quark, $t \to Wb \to jjb$ and the charged Higgs decay to τ-leptons.

Table 1: The parameters used in the current analysis of the signal with the corresponding polarisation asymmetry. In general, H^- would decay to τ_L^- and τ_R^-, $H^- \to \tau_R^- \bar{\nu} + \tau_L^- \psi$, depending on the asymmetry. For the purely decay $H^- \to \tau_R^- \bar{\nu}$ (as in MSSM), the asymmetry is -1. The signal to be studied is $H^- \to \tau_L^- \psi$.

	M_* (TeV)	δ_ν, δ	$m_{H^\pm}$ (GeV)	$\tan\beta$	A_{LR}	m_ν (eV)
Sig.-1	2	4, 4	219.9	30	~ 1	$0.5\ 10^{-3}$
Sig.-2	20	3, 3	365.4	45	~ 1	0.05
Sig.-3	1	5, 6	506.2	4	~ 1	0.05
Sig.-4	100	6, 6	250.2	35	~ -1	0.005
Sig.-5	10	4, 5	350.0	20	~ -1	0.04
Sig.-6	50	5, 5	450.0	25	~ -1	0.04

The major backgrounds are the single top production $gb \to Wt$, and $t\bar{t}$ production with one $W^+ \to jj$ and the other $W^- \to \tau_L^- \bar{\nu}$. Depending on the polarisation asymmetry, $H^- \to \tau_R^- \bar{\nu}$ will contribute as an additional background. In Table 2, we list the rates for the signal and for the backgrounds.

Table 2: The expected rates ($\sigma\times$ BR), for the signal $gb \to tH^\pm$ with $H^- \to \tau_R^- \bar{\nu} + \tau_L^- \psi$ and $t \to jjb$, and for the backgrounds: Wt and $t\bar{t}$ with $W^- \to \tau_L^- \bar{\nu}$ and $W^+ \to jj$. We assume an inclusive $t\bar{t}$ production cross section of 590 pb. In the last columns, we compare the $H^- \to \tau_R^- \bar{\nu}$ branching ratios in this model to the corresponding MSSM branching ratios.

Process	$\sigma \times$ BR (pb)	BR	BR(MSSM)
Signal-1	1.56	0.73	0.37
Signal-2	0.15	1.0	0.15
Signal-3	0.04	1.0	0.01
$t\bar{t}$	84.11		
Wtb ($p_T > 30$ GeV)	47.56		

Depending on the parameters M_*, m_D, δ, $m_{H^\pm}$ and $\tan\beta$, the $\tau\nu$ decay of the charged Higgs can be enhanced or suppressed compared to the MSSM case. In Figure 1, we show how the other decays of the charged Higgs are affected in this framework; for the chosen values of M_* and δ, the decay branchings are similar to MSSM for small values of m_D while at larger m_D, the $\tau\nu$ decay mode becomes strongly enhanced.

The reconstruction of the transverse mass is not enough to distinguish between the MSSM and the singlet neutrinos in large extra dimensions. The differences in these two scenarii are best seen in the distribution of $p^\pi/E^{\tau-jet}$, the fraction of the energy carried by the charged track.



Figure 1: Charged Higgs decays in models with a singlet neutrino in large extra dimensions for $M_* = 2 \times 10^4$ GeV, $\delta = 3$ and $\tan\beta = 45$. For small values of m_D, we see similar decay branchings as in MSSM. As m_D gets larger, $H^{\pm} \to \tau\nu$ becomes dominant below and above the top-quark mass.



Figure 2: The distribution of the ratio of the charged pion track momentum in one prong τ decay to the τ-jet energy for $m_A = 350$ GeV, $\tan\beta = 45$, $M_* = 20$ TeV, $\delta = 3$ and $m_\nu = 0.05$ eV. In the 2HDM-II, this ratio would peak near 0 and 1 as shown while in other models, the actual distribution of this ratio would depend on the polarization asymmetry since both left and right handed τ's would contribute. In the case shown, the asymmetry is ~ 1 and the ratio peaks near the center of the distribution.

3 Conclusions

Large extra dimensions models with TeV scale quantum gravity postulate the existence of additional dimensions where gravity (and possibly other fields) propagate. The size of the extra dimensions are constrained to the sub-millimeter level since no experimental deviations from the Newtonian gravity has been observed at distances larger than ~ 0.2 millimeter. The right handed neutrino can freely propagate into the extra dimensions because it has no quantum numbers to constrain it to the SM brane. The interactions between the bulk neutrino and the SM fields on the brane can generate Dirac neutrino masses consistent with the atmospheric neutrino oscillations without implementing the seesaw mechanism. There are no additional Higgs bosons required in these models. The charged Higgs productions are therefore the same as in the two Higgs doublet models. The charged Higgs can decay to both the right and the left handed τ-leptons, $H^- \to \tau_R^- \bar{\nu} + \tau_L^- \psi$ whereas in the 2HDM-II such as MSSM, only the right handed τ decay of the H^- is possible through the τ Yukawa coupling: $H^- \to \tau_R^- \bar{\nu}$. The τ decay of the charged Higgs has been studied in details for ATLAS and CMS. In the current study, we focus on the decay $H^- \to \tau_L^- \psi$ at the LHC for Higgs masses larger than the top-quark mass.

Although the observation of a signal in the transverse mass distribution can be used to claim discovery of the charged Higgs, it is insufficient to select the scenario that is realized. Additionally, by reconstructing the fraction of the energy carried by the charged track in the one-prong τ decay, it is possible to distinguish whether the scenario is the ordinary two Higgs doublet model or not. The further measurement of the polarisation asymmetry might provide a distinctive evidence for models with singlet neutrino in large extra dimensions.

Acknowledgements

I would like to thank K. A. Assamagan and Y. Coadou for fruitful collaborations.

References

1. C. D. Hoyle, U. Schmidt, B. R. Heckel, E. G. Adelberger, J. H. Gundlach, D. J. Kapner and H. E. Swanson, Phys. Rev. Lett. **86**, 1418 (2001) [arXiv:hep-ph/0011014].
2. K. Agashe, N. G. Deshpande and G. H. Wu, Phys. Lett. B **489**, 367 (2000) [arXiv:hep-ph/0006122];
3. K.A. Assamagan and Y. Coadou, ATLAS Internal Note, ATL-PHYS-2000-031 (2000); K. A. Assamagan, Acta Phys. Polon. B **31**, 863 (2000); K. A. Assamagan and Y. Coadou, Acta Phys. Polon. B **33** 707 (2002); K. A. Assamagan, Y. Coadou and A. Deandrea, EPJdirect (Vol. 4) **C9** 1 (2002) [arXiv:hep-ph/0203121].
4. R. Kinnunen, "The $H^{\pm} \to \tau\nu$ mode in CMS" and "Signatures of Heavy Charged Higgs Bosons at the LHC", in Les Houches Workshop (1999), "The Higgs working group: Summary report," arXiv:hep-ph/0002258; D. Cavalli *et al.*, "The Higgs working group: Summary report," in Les Houches Workshop (2001), arXiv:hep-ph/0203056.
5. K. A. Assamagan and A. Deandrea, Phys. Rev. D **65** (2002) 076006 [arXiv:hep-ph/0111256].